
Attal / Joye / Pillet

Open Quantum Systems I

The Hamiltonian Approach

This is the ?rst in a series of three volumes dedicated to the lecture notes of the Summer School ”Open Quantum Systems” which took place at the Institut Fourier in Grenoble from June 16th to July 4th 2003. The contributions presented in these volumes are revised and expanded versions of the notes provided to the students during the School. Closed vs. Open Systems By de?nition, the time evolution of a closed physical systemS is deterministic. It is usually described by a differential equation x ? = X(x) on the manifold M of t t possible con?gurations of the system. If the initial con?guration x ? M is known 0 then the solution of the corresponding initial value problem yields the con?guration x at any future time t. This applies to classical as well as to quantum systems. In the t classical case M is the phase space of the system and x describes the positions and t velocities of the various components (or degrees of freedom) ofS at time t. Inthe quantum case, according to the orthodoxinterpretation of quantum mechanics, M is a Hilbert space and x a unit vector – the wave function – describing the quantum t state of the system at time t. In both cases the knowledge of the state x allows t to predict the result of any measurement made onS at time t.

This is the ?rst in a series of three volumes dedicated to the lecture notes of the Summer School "Open Quantum Systems" which took place at the Institut Fourier in Grenoble from June 16th to July 4th 2003. The contributions presented in these volumes are revised and expanded versions of the notes provided to the students during the School. Closed vs. Open Systems By de?nition, the time evolution of a closed physical systemS is deterministic. It is usually described by a differential equation x ? = X(x) on the manifold M of t t possible con?gurations of the system. If the initial con?guration x ? M is known 0 then the solution of the corresponding initial value problem yields the con?guration x at any future time t. This applies to classical as well as to quantum systems. In the t classical case M is the phase space of the system and x describes the positions and t velocities of the various components (or degrees of freedom) ofS at time t. Inthe quantum case, according to the orthodoxinterpretation of quantum mechanics, M is a Hilbert space and x a unit vector - the wave function - describing the quantum t state of the system at time t. In both cases the knowledge of the state x allows t to predict the result of any measurement made onS at time t.



69,54 €
64,99 € (zzgl. MwSt.)

Lieferfrist: bis zu 10 Tage

Artikelnummer: 9783540309918
Medium: Buch
ISBN: 978-3-540-30991-8
Verlag: Springer
Erscheinungstermin: 07.06.2006
Sprache(n): Englisch
Auflage: 1. Auflage 2006
Serie: Lecture Notes in Mathematics
Produktform: Kartoniert
Gewicht: 534 g
Seiten: 329
Format (B x H): 155 x 235 mm

Kundenservice Verlag Dr. Otto Schmidt KG
Neumannstraße 10, 40235 Düsseldorf | kundenservice@otto-schmidt.de | 0221-93738-997 (Inland)