


Applied Finite Group Actions

Also the present second edition of this book is an introduction to the theory of classification, enumeration, construction and generation of finite unlabeled structures in mathematics and sciences. Since the publication of the first edition in 1991 the constructive theory of unlabeled finite structures has made remarkable progress. For example, the first designs with moderate parameters were constructed, in Bayreuth, by the end of 1994 ([9]). The crucial steps were - the prescription of a suitable group of automorphisms, i. e. a stabilizer, and the corresponding use of Kramer-Mesner matrices, together with - an implementation of an improved version of the LLL-algorithm that allowed to find 0-1-solutions of a system of linear equations with the Kramer-Mesner matrix as its matrix of coefficients. The Kramer-Mesner matrices can be considered as submatrices form A " (see the chapter on group actions on posets, semigroups and lattices). They are associated with the action of the prescribed group G which is a permutation group on a set X of points induced on the power set of X . Hence the discovery of the first 7-designs with small parameters is due to an application of finite group actions. This method used by A. Betten, R. Laue, A. Wassermann and the present author is described in a section that was added to the manuscript of the first edition.

Also the present second edition of this book is an introduction to the theory of classification, enumeration, construction and generation of finite unlabeled structures in mathematics and sciences. Since the publication of the first edition in 1991 the constructive theory of unlabeled finite structures has made remarkable progress. For example, the first designs with moderate parameters were constructed, in Bayreuth, by the end of 1994 ([9]). The crucial steps were - the prescription of a suitable group of automorphisms, i. e. a stabilizer, and the corresponding use of Kramer-Mesner matrices, together with - an implementation of an improved version of the LLL-algorithm that allowed to find 0-1-solutions of a system of linear equations with the Kramer-Mesner matrix as its matrix of coefficients. The Kramer-Mesner matrices can be considered as submatrices form A " (see the chapter on group actions on posets, semigroups and lattices). They are associated with the action of the prescribed group G which is a permutation group on a set X of points induced on the power set of X . Hence the discovery of the first 7-designs with small parameters is due to an application of finite group actions. This method used by A. Betten, R. Laue, A. Wassermann and the present author is described in a section that was added to the manuscript of the first edition.



160,49 €

149,99 € (zzgl. MwSt.)

Lieferfrist: bis zu 10 Tage

Artikelnummer: 9783540659419

Medium: Buch

ISBN: 978-3-540-65941-9

Verlag: Springer Netherlands

Erscheinungstermin: 18.08.1999

Sprache(n): Englisch

Auflage: 2. Auflage 1999

Serie: Algorithms and Combinatorics

Produktform: Gebunden

Gewicht: 1880 g

Seiten: 454

Format (B x H): 155 x 235 mm

