


Braids and Self-Distributivity

The aim of this book is to present recently discovered connections between Artin's braid groups E_n and left self-distributive systems (also called LD systems), which are sets equipped with a binary operation satisfying the left self-distributivity identity $x(yz) = (xy)(xz)$. (LD) Such connections appeared in set theory in the 1980s and led to the discovery in 1991 of a left invariant linear order on the braid groups. Braids and self-distributivity have been studied for a long time. Braid groups were introduced in the 1930s by E. Artin, and they have played an increasing role in mathematics in view of their connection with many fields, such as knot theory, algebraic combinatorics, quantum groups and the Yang-Baxter equation, etc. LD-systems have also been considered for several decades: early examples are mentioned in the beginning of the 20th century, and the first general results can be traced back to Belousov in the 1960s. The existence of a connection between braids and left self-distributivity has been observed and used in low dimensional topology for more than twenty years, in particular in work by Joyce, Brieskorn, Kauffman and their students. Brieskorn mentions that the connection is already implicit in (Hurwitz 1891). The results we shall concentrate on here rely on a new approach developed in the late 1980s and originating from set theory.

The aim of this book is to present recently discovered connections between Artin's braid groups E_n and left self-distributive systems (also called LD systems), which are sets equipped with a binary operation satisfying the left self-distributivity identity $x(yz) = (xy)(xz)$. (LD) Such connections appeared in set theory in the 1980s and led to the discovery in 1991 of a left invariant linear order on the braid groups. Braids and self-distributivity have been studied for a long time. Braid groups were introduced in the 1930s by E. Artin, and they have played an increasing role in mathematics in view of their connection with many fields, such as knot theory, algebraic combinatorics, quantum groups and the Yang-Baxter equation, etc. LD-systems have also been considered for several decades: early examples are mentioned in the beginning of the 20th century, and the first general results can be traced back to Belousov in the 1960s. The existence of a connection between braids and left self-distributivity has been observed and used in low dimensional topology for more than twenty years, in particular in work by Joyce, Brieskorn, Kauffman and their students. Brieskorn mentions that the connection is already implicit in (Hurwitz 1891). The results we shall concentrate on here rely on a new approach developed in the late 1980s and originating from set theory.



106,99 €
99,99 € (zzgl. MwSt.)

Lieferfrist: bis zu 10 Tage

Artikelnummer: 9783764363437
Medium: Buch
ISBN: 978-3-7643-6343-7
Verlag: Springer
Erscheinungstermin: 01.07.2000
Sprache(n): Englisch
Auflage: 1. Auflage 2000
Serie: Progress in Mathematics
Produktform: Gebunden
Gewicht: 1133 g
Seiten: 623
Format (B x H): 160 x 241 mm

